

Table 2 Body weight (gm) (mean \pm SE) of Nile tilapia fingerlings in different ponds

Pond	Experimental period (days)								
	0	15	30	45	60	75	90	105	120
P1	5.7 \pm	5.9 \pm	6.8 \pm	8.9 \pm	9.1 \pm	11.3 \pm	13.6 \pm	15.7 \pm	18.2 \pm
	0.02 ^a	0.02 ^c	0.03 ^c	0.04 ^c	0.03 ^c	0.03 ^c	0.02 ^c	0.02 ^c	0.02 ^c
P2	5.7 \pm	7.1 \pm	11.2 \pm	15.9 \pm	21.2 \pm	27.2 \pm	33.1 \pm	42.3 \pm	52.4 \pm
	0.02 ^a	0.04 ^b	0.03 ^b	0.04 ^b	0.05 ^b	0.05 ^b	0.05 ^b	0.53 ^b	0.57 ^b
P3	5.7 \pm	8.7 \pm	12.2 \pm	17.1 \pm	22.8 \pm	30.3 \pm	39.8 \pm	51.5 \pm	63.7 \pm
	0.02 ^a	0.08 ^a	0.06 ^a	0.12 ^a	0.13 ^a	0.27 ^a	0.22 ^a	0.27 ^a	0.50 ^a

Note: Means with the same letter in each column are not significantly different ($P>0.05$)