

Table 3 Truss network of distance ratio of *Polydactylus sextarius*, *Nemipterus japonicus* and *Pampus argenteus*

Character (×100%)	<i>Polydactylus sextarius</i>	<i>Nemipterus japonicus</i>	<i>Pampus argenteus</i>
2/1	0.32 ±0.01	0.34 ±0.01	0.67 ±0.03
3/1	0.30 ±0.01	0.31 ±0.02	0.25 ±0.01
4/3	0.80 ±0.05	0.93 ±0.06	0.36 ±0.02
6/3	0.21 ±0.03	0.30 ±0.02	0.04 ±0.00
7/1	0.18 ±0.02	0.15 ±0.01	0.08 ±0.01
8/1	0.15 ±0.01	0.11 ±0.00	0.09 ±0.00
9/1	0.20 ±0.01	0.10 ±0.00	0.29 ±0.02
10/1	0.22 ±0.01	0.30 ±0.02	0.34 ±0.02
11/1	0.16 ±0.01	0.25 ±0.02	——
13/1	0.17 ±0.01	0.12 ±0.01	0.42 ±0.04
14/1	0.17 ±0.01	——	——
BD/1	0.22 ±0.01	0.32 ±0.01	0.21 ±0.01
AD/1	0.20 ±0.01	0.16 ±0.02	0.23 ±0.02
EF1/1	0.45 ±0.02	0.46 ±0.03	0.63 ±0.02
EF2/1	0.33 ±0.01	——	——
DF1/1	0.19 ±0.01	0.09 ±0.00	0.40 ±0.04
DF2/1	0.49 ±0.01	——	——
CE/1	0.26 ±0.01	0.30 ±0.02	——
CD/1	0.33 ±0.01	0.33 ±0.02	——
F1H1/1	0.14 ±0.00	0.55 ±0.04	0.50 ±0.02
F2H2/1	0.21 ±0.02	——	——
IJ/1	0.15 ±0.00	0.11 ±0.00	0.10 ±0.00
GJ/1	0.24 ±0.02	0.21 ±0.01	0.13 ±0.00
GH1/1	0.43 ±0.02	0.17 ±0.01	0.10 ±0.00
GH2/1	0.19 ±0.01	——	——
AC/1	0.14 ±0.01	0.17 ±0.01	——
H1I/1	0.58 ±0.02	0.18 ±0.02	0.14 ±0.01
H2I/1	0.27 ±0.02	——	——
EH1/1	0.37 ±0.02	0.31 ±0.01	0.67 ±0.02
EH2/1	0.30 ±0.02	——	——
DE/1	0.54 ±0.01	0.51 ±0.01	——
BC/1	0.39 ±0.01	0.12 ±0.02	——
H1J/1	0.54 ±0.02	——	0.10 ±0.01
H2J/1	0.21 ±0.03	——	——
F1G/1	0.55 ±0.03	0.58 ±0.01	0.58 ±0.02
F2G/1	0.33 ±0.04	——	——
GI/1	0.18 ±0.02	0.16 ±0.00	0.10 ±0.01
AB/1	0.27 ±0.01	0.34 ±0.02	0.58 ±0.02
CF1/1	0.33 ±0.03	0.33 ±0.02	——
CF2/1	0.39 ±0.03	——	——

Note: Where: *Polydactylus sextarius* have two dorsal fin, *Pampus argenteus* have no pelvic fin, and ‘——’ represent empty, (1) body length, (2) height, (3) head length, (4) head height, (5) snout length, (6) eye diameter, (7) caudal peduncle length, (8) caudal peduncle height, (9) the first dorsal fin length(or dorsal fin length), (10) pectoral fin length, (11) pelvic fin length, (12) anal fin base length, (13) anal fin length, and (14) the second dorsal length (Fig 1)